

Quantification of loss of heterozygosity in Fischer p53 RRRC knockout rats using ddPCR

JR E Sp

<u>Jiajie Jessica Xu^{1,2}</u>, Sarah A. Hansen¹, Susheel B. Busi¹, Taybor W. Parker¹, Marcia L. Hart¹, Elizabeth Bryda^{1,3}, James M. Amos-Landgraf^{1,3}

¹Department of Veterinary Pathobiology, University of Missouri, Columbia, MO,

²College of Veterinary Medicine, Texas A&M University, College Station, TX, ³Rat Resource and Research Center, Columbia MO

Hypothesis: Fischer p53+/- knockout rats develop tumors through loss of heterozygosity (LOH), and the mechanism of LOH can be determined by accurately quantifying gene copy number through the use of droplet digital PCR (ddPCR).




Figure 1: Knudson's 2 hit hypothesis for cancer development, with 4 potential mechanisms for LOH.

p53 knockout rats:

- germline competent rat ES cells
- variable phenotype between strains (fig 2)
- better models for some cancers than mice (fig 3).






Figure 2: Tumor variation in p53+/+ and p53+/- Sprague Dawley and Fischer rats



Figure 3: Radiograph (A) and gross (B) depiction of osteosarcoma from p53 knockout rat. Histology from bone of rat (C) and mouse (D). Like humans, rats form true lamellae (L), which mice lack.

Materials and methods





Control Samples

■GFP ■p53

Summary

- 1. Developed LOH ddPCR assay.
- 2. Demonstrated LOH in Fischer p53^{+/-} rats, consistent with other cancer models.
- 3. ddPCR has a increased sensitivity compared to qPCR, but further refinement is needed for accurate quantification of copy number.

Acknowledgements

assay necessary for accurate

gene number quantification.

Student funding:



